

Diagnostic criteria for drug rash and eosinophilia with systemic symptoms

Dear Editor,

We found the paper by Jmeian *et al.* concerning a case of allopurinol-related drug rash with eosinophilia and systemic symptoms (DRESS) in a patient of chronic kidney disease and gout, interesting.^[1] The presence of eosinophilia, the temporal relationship of the symptoms with the initiation of treatment with allopurinol, and the marked improvement on withdrawal of the drug along with the administration of systemic corticosteroids were features suggesting the diagnosis of DRESS.

DRESS is considered a severe drug reaction with a case fatality rate of 10–20%.^[2] The diagnosis is sometimes difficult since the clinical manifestations may be incomplete or nonspecific, and it can also present as a purely systemic disease without any cutaneous involvement.^[3,4]

Multiple diagnostic criteria have been developed and used to standardize the diagnosis and management of DRESS, albeit with limited success.

Bocquet *et al.* were the first who proposed criteria for DRESS [Table 1].^[5] The Registry of Severe Cutaneous Adverse Reaction (RegiSCAR) group suggested criteria for hospitalized patients with a drug rash to diagnose DRESS syndrome [Table 2].^[6] RegiSCAR constitutes a European RegiSCAR, including Stevens–Johnson syndrome, toxic epidermal necrolysis, acute generalized exanthematous pustulosis, and DRESS. In an effort to define more accurately the DRESS syndrome, a scoring system has also been developed the RegiSCAR scoring system [Table 3].^[6] A Japanese group suggested another set of diagnostic criteria; however, universal adaptation of this criteria may be limited since one of the criteria includes HHV-6 activation and some tests, such as measurement of IgG titer anti-HHV 6, are yet not routinely available [Table 4].^[7]

DRESS is a challenging drug adverse reaction which can cause life-threatening organ dysfunction. Clinicians must be alert to this possibility to reach the correct diagnosis and institute the appropriate management.

Financial support and sponsorship

Nil.

Table 1: Bocquet *et al.* proposed criteria for diagnosis of drug rash and eosinophilia with systemic symptoms/ drug-induced hypersensitivity

DRESS is confirmed by the presence of 1 and 2 and 3

Cutaneous drug eruption

Adenopathy >2 cm in diameter or hepatitis (liver transaminases >2 times of normal or interstitial nephritis or interstitial pneumonia or carditis

Hematologic abnormalities eosinophilia $>1.5 \times 10^9/L$ or atypical lymphocytes

DRESS: Drug rash and eosinophilia with systemic symptoms

Table 2: Registry of severe cutaneous adverse reaction criteria for diagnosis of drug rash and eosinophilia with systemic symptoms

1. Hospitalization*
2. Reaction suspected to be drug-related*
3. Acute rash*
4. Fever $>38^\circ C^\dagger$
5. Enlarged lymph nodes at a minimum of 2 sites[†]
6. Involvement of at least 1 internal organ[†]
7. Blood count abnormalities[†]
 - Lymphocytes above or below normal limits
 - Eosinophils above the laboratory limits
 - Platelets below the laboratory limits

*Necessary criteria are required for making the diagnosis. [†]Three out of four criteria are required

Table 3: Registry of severe cutaneous adverse reaction diagnosis score for drug rash and eosinophilia with systemic symptoms

Features	No	Yes	Unknown
Fever $>38.5^\circ C$	–1	0	–1
Enlarged lymph nodes (>2 sites, >1 cm)	0	1	0
Atypical lymphocytes	0	1	0
Eosinophilia			0
700–1499 or 10%–19.9%	0	1	
≥ 1500 or $\geq 20\%$		2	
Skin rash			
Extent $>50\%$	0	1	
At least 2: Edema, infiltration, purpura, scaling	–1	1	
Biopsy suggesting DRESS	–1	0	
Internal organ involvement			
One	0	1	0
Two or more		2	
Resolution in ≥ 15 days	–1	0	–1
Evaluation of other potential causes (antinuclear antibody, blood culture, serology for HAV/ HBV/ HCV, chlamydia/ mycoplasma)	0	1	0
If none of these positive and >3 are negative			

Conflicts of interest

There are no conflicts of interest.

Table 4: Japanese group's criteria for diagnosis of drug rash and eosinophilia with systemic symptoms/ drug-induced hypersensitivity

1. Maculopapular rash developing >3 weeks after starting with the suspected drug
2. Prolonged clinical symptoms 2 weeks after discontinuation of the suspected drug
3. Fever >38°C
4. Liver abnormalities (alanine aminotransferase >100U/L)*
5. Leucocyte abnormalities (at least one present)
 - (a) Leucocytosis (>11×10⁹/L)
 - (b) Atypical lymphocytosis (>5%)
 - (c) Eosinophilia (>1.5×10⁹/L)
6. Lymphadenopathy
7. Human Herpes 6 reactivation

The diagnosis is confirmed by the presence of the 7 criteria (typical DIHS) or of the five (1-5) (atypical DIHS). *This can be replaced by other organ involvement, such as renal involvement

Ashok Kumar Pannu¹, Atul Saroch²

¹Department of Internal Medicine, Post Graduate Institute of Medical Education and Research, Chandigarh, ²Department of Medicine, Post Graduate Institute of Medical Education and Research, Dr. RML Hospital, New Delhi, India

Address for correspondence: Dr. Ashok Kumar Pannu, Department of Internal Medicine, Post Graduate Institute of Medical Education and Research, 4th Floor, F Block, Chandigarh - 160 012, India.
E-mail: gawaribacchi@gmail.com

References

1. Jmeian A, Hawatmeh A, Shamoan R, Shamoan F, Guma M. Skin rash, eosinophilia, and renal impairment in a patient recently started on allopurinol. *J Family Med Prim Care* 2016;5:479-81.
2. Kumari R, Timshina DK, Thappa DM. Drug hypersensitivity syndrome. *Indian J Dermatol Venereol Leprol* 2011;77:7-15.
3. Sasidharanpillai S, Riyaz N, Rajan U, Binitha MP, Khader A, Mariyath OK, *et al.* Drug reaction with eosinophilia and systemic symptoms: Observations from a tertiary care institution. *Indian J Dermatol Venereol Leprol* 2014;80:221-8.
4. Ben m'rad M, Leclerc-Mercier S, Blanche P, Franck N, Rozenberg F, Fulla Y, *et al.* Drug-induced hypersensitivity syndrome: Clinical and biologic disease patterns in 24 patients. *Medicine (Baltimore)* 2009;88:131-40.
5. Bocquet H, Bagot M, Roujeau JC. Drug-induced pseudolymphoma and drug hypersensitivity syndrome (Drug Rash with Eosinophilia and Systemic Symptoms: DRESS). *Semin Cutan Med Surg* 1996;15:250-7.
6. Kardaun SH, Sidoroff A, Valeyrie-Allanore L, Halevy S, Davidovici BB, Mockenhaupt M, *et al.* Variability in the clinical pattern of cutaneous side-effects of drugs with systemic symptoms: Does a DRESS syndrome really exist? *Br J Dermatol* 2007;156:609-11.
7. Shiohara T, Iijima M, Ikezawa Z, Hashimoto K. The diagnosis of a DRESS syndrome has been sufficiently established on the basis of typical clinical features and viral reactivations. *Br J Dermatol* 2007;156:1083-4.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:



Website:
www.jfmpc.com

DOI:
10.4103/2249-4863.222050

How to cite this article: Pannu AK, Saroch A. Diagnostic criteria for drug rash and eosinophilia with systemic symptoms. *J Family Med Prim Care* 2017;6:693-4.

© 2017 Journal of Family Medicine and Primary Care | Published by Wolters Kluwer - Medknow